

Mr Gavin A J Morrison MA FRCS
Consultant Otolaryngologist
Guy's & St. Thomas' Hospitals

73 Harley Street
London W1G 8QJ

Tel: 020 7487 4446
Fax: 020 7224 6381

email: gajm@gavinmorrison.com

The Lister Hospital
020 7730 7733

The London Clinic
020 7935 4444

Harley Street Clinic
020 7935 7700

The Portland Hospital
020 7580 4400

INFORMATION FOR PATIENTS ON HOSPITAL DISCHARGE
Following Ear Surgery – Stapedectomy

- 1. EAR PACK:** You will be discharged with an antiseptic ribbon pack (BIPP) already in the ear canal. This must be left in place until you return to the consulting room for Mr Morrison to remove it, usually after 7 - 14 days.
- 2. KEEP EAR DRY:** Please keep the ear dry. Do not let water get in the ear for two weeks. You can wash your hair, ideally with someone to help, after a couple of days. When doing this hold a towel over the ear or hold a glass or cup over the ear to protect it from water. This way you can use a hand-held shower. Do not put your whole head under water in the shower or bath. Dry your hair with a hair dryer, and avoid rubbing behind the ear.

You will need to keep water out of the ear canal for longer, until it has all healed fully, which often takes about a month.

- 3. NOSE BLOWING:** You should not put any pressure on the Eustachian tube or ear. This means you should not blow your nose at all for 1 – 2 weeks, and then not at all vigorously.

- 4. MEDICATIONS:**
Antibiotics

If a course of antibiotics has been prescribed, please take these as indicated on the bottle.

Pain Killers

Simple analgesics such as Paracetamol, or Nurofen (Ibuprofen) may be sufficient. Unless you are allergic (or have asthma with Aspirin/Nurofen sensitivity), you can take both these pain killers at the same time to increase their effect. Take them as directed on the packet 3 –4 times per day if necessary. You can use your own supply of these, bought from the chemist.

You may have been given more powerful pain killers such as Co-Codamol, Tylex, or Codeine Phosphate. Use these if necessary but do not take Tylex or Co-Codamol at the same time as Paracetamol. Voltarol (Diclofenac) may have been prescribed on discharge. This is like Ibuprofen (Nurofen), and so should not be taken at the same time as Voltarol.

/continued

5. **HEARING LOSS – BLOCKED EAR:** The operated ear will feel very blocked, because of the pack and also because the ear will be bruised and full of blood after the surgery. This will all take some weeks to gradually clear. Expect the ear to feel very deaf to begin with. After a week or more the middle ear may start to clear, and sudden loud crackling noises, popping or other types of tinnitus, (ringing or buzzing) are normal and will clear.
6. **DIZZINESS:** Some mild dizziness is to be expected at first, with unsteadiness when walking or moving the head suddenly. This should gradually improve over a few days or a few weeks. If very troublesome, you might need to take an anti-dizzy medication such as Stugeron, 15 mg every 8 hours. This may be bought over the counter at a chemist.
7. **FOLLOW UP:** Your first out-patient clinic visit should usually be arranged at about 7 days from the operation. Please telephone the office to book it at your convenience. At this visit Mr Morrison will be able to remove the ear pack.
8. **BLEEDING:** It is normal to have a yellow or blood stained secretion from the ear canal for up to a week after the surgery. You can replace the cotton wool in the ear when necessary. Do not remove the BIPP pack, beyond it.

Bleeding with bulging behind the ear might indicate a haematoma. You should phone Mr Morrison's secretary.
9. **SICK LEAVE:** You are usually advised to take 2 - 3 weeks sick leave from work if this is possible following your operation. If not, try to return to work on an easy schedule, as stress should be avoided.
10. **EXERCISE:** Please avoid any strenuous exercise in the first 2-3 weeks after the operation. Brisk walking & everyday activities are fine but swimming, gym-work or sports are best avoided.
11. **DRIVING:** You can drive after a few days or a week, so long as you feel well, with no dizziness. If in doubt, wait for a bit longer.
12. **OTHER SYMPTOMS:** The outer ear may feel numb for a month or so, until the skin nerves which had to be cut have grown back. There may be an altered, metallic or tingling taste in the tongue, after the surgery. This usually settles, but can take a few months.
13. **EMERGENCIES:** If you need to contact Mr Morrison for general advice or in an emergency firstly please try the office telephone number 0207 487 4446.

If there is no response on the above number and you need to contact Mr Morrison out of hours please telephone the Hospital & ward where you stayed for your operation. The Nurses will be able to get a message to Mr Morrison at home or on his mobile phone so that he can advise. Similarly, the other hospitals listed on the letterhead could try to contact him for you. If Mr Morrison is going to be away, he will usually have suggested an ENT Consultant Colleague who might cover. Finally, NHS emergency care is also available from Guy's & St Thomas' Hospitals, or from your nearest NHS hospital with an ENT Department.